

Appl. No. 10/526,822
Reply to Office Action of January 7, 2008

RECEIVED
CENTRAL FAX CENTER

APR 07 2008

Amendments to the Claims:

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1. (Currently amended) A clear ophthalmic solution comprising latanoprost having a concentration of 0.005 to 0.01% (W/V) as an active ingredient and benzalkonium chloride as a preservative wherein white turbidity due to a change of formulation is prevented by at least one means selected from the following 1) to 2);

1) using 0.003 to 0.01% (W/V) benzalkonium chloride represented by the formula of $[C_6H_5CH_2N(CH_3)_2R]Cl$ (wherein R is alkyl having 12 carbon atoms) as the preservative and

2) adding a tonicity agent consisting essentially of a nonionic tonicity agent.

2-3. (Cancelled).

RECEIVED
CENTRAL FAX CENTER

APR 07 2008

Appl. No. 10/526,822

Reply to Office Action of January 7, 2008

4. **(Previously presented)** The ophthalmic solution as claimed in claim 9, wherein the nonionic tonicity agent is used and is selected from the group consisting of glycerin, mannitol, polyethylene glycol, propylene glycol, trehalose or sucrose.

5. **(Currently amended)** A method of preventing white turbidity due to a change of formulation in an ophthalmic solution comprising latanoprost having a concentration of 0.005 to 0.01% (W/V) as an active ingredient and benzalkonium chloride as a preservative by at least one means selected from the following 1) to 2):

1) using 0.003 to 0.01% (W/V) benzalkonium chloride represented by the formula of $[C_6H_5CH_2N(CH_3)_2R] Cl$ (wherein R is alkyl having 12 carbon atoms) as the preservative and

2) adding a tonicity agent consisting essentially of a nonionic tonicity agent.

6-7. **(Cancelled).**

Appl. No. 10/526,822

Reply to Office Action of January 7, 2008

8. **(Previously presented)** The method of preventing white turbidity as claimed in claim 10, wherein the nonionic tonicity agent is used and is selected from the group consisting of glycerin, mannitol, polyethylene glycol, propylene glycol, trehalose or sucrose.

9. **(Original)** The ophthalmic solution as claimed in claim 2, wherein a latanoprost concentration is 0.005% (W/V).

10. **(Original)** The method of preventing white turbidity as claimed in claim 6, wherein a latanoprost concentration is 0.005% (W/V).